

B.06) Combinations and Permutations

<http://www.mathsisfun.com/combinatorics/combinations-permutations.html>

We use the word "combination" loosely, without thinking if the **order** of things is important. In other words:



"My fruit salad is a combination of apples, grapes and bananas" We don't care what order the fruits are in, they could also be "bananas, grapes and apples" or "grapes, apples and bananas", its the same fruit salad.



"The combination to the safe was 472". Now we **do** care about the order. "724" would not work, nor would "247". It has to be exactly **4-7-2**.

So, in Mathematics we use more *precise* language:

- If the order doesn't matter, it is a **Combination**.
- If the order **does** matter it is a **Permutation**.



So, we should really call this a "Permutation Lock"!

In other words: a Permutation is an **ordered** Combination.

Permutations

There are basically two types of permutation:

1. **Repetition is Allowed**: such as the lock above. It could be "333".
2. **No Repetition**: for example the first three people in a running race. You can't be first *and* second.

1. Permutations with Repetition

These are the easiest to calculate.

When you have n things to choose from ... you have n choices each time!

When choosing r of them, the permutations are:

$$n \times n \times \dots (r \text{ times})$$

(In other words, there are n possibilities for the first choice, THEN there are n possibilities for the second choice, and so on, multiplying each time.)

Which is easier to write down using an [exponent](#) of r :

$$n \times n \times \dots (r \text{ times}) = n^r$$

Example: in the lock above, there are 10 numbers to choose from (0,1,..9) and you choose 3 of them:

$$10 \times 10 \times \dots (3 \text{ times}) = 10^3 = 1,000 \text{ permutations}$$

So, the formula is simply:

$$n^r$$

where n is the number of things to choose from, and you choose r of them
(Repetition allowed, order matters)

2. Permutations without Repetition

In this case, you have to **reduce** the number of available choices each time.



For example, what order could 16 pool balls be in?

After choosing, say, number "14" you can't choose it again.

So, your first choice would have 16 possibilities, and your next choice would then have 15 possibilities, then 14, 13, etc. And the total permutations would be:

$$16 \times 15 \times 14 \times 13 \times \dots = 20,922,789,888,000$$

But maybe you don't want to choose them all, just 3 of them, so that would be only:

$$16 \times 15 \times 14 = 3,360$$

In other words, there are 3,360 different ways that 3 pool balls could be selected out of 16 balls.

But how do we write that mathematically? Answer: we use the "[factorial function](#)"



The **factorial function** (symbol: !) just means to multiply a series of descending natural numbers. Examples:

- $4! = 4 \times 3 \times 2 \times 1 = 24$
- $7! = 7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1 = 5,040$
- $1! = 1$

Note: it is generally agreed that $0! = 1$. It may seem funny that multiplying no numbers together gets you 1, but it helps simplify a lot of equations.

So, if you wanted to select **all** of the billiard balls the permutations would be:

$$16! = 20,922,789,888,000$$

But if you wanted to select just 3, then you have to stop the multiplying after 14. How do you do that? There is a neat trick ... you divide by $13!$...

$$\frac{16 \times 15 \times 14 \times 13 \times 12 \dots}{13 \times 12 \dots} = 16 \times 15 \times 14 = 3,360$$

Do you see? $16! / 13! = 16 \times 15 \times 14$

The formula is written:

$$\frac{n!}{(n - r)!}$$

where n is the number of things to choose from, and you choose r of them
(No repetition, order matters)

Examples:

Our "order of 3 out of 16 pool balls example" would be:

$$\frac{16!}{(16-3)!} = \frac{16!}{13!} = \frac{20,922,789,888,000}{6,227,020,800} = 3,360$$

(which is just the same as: $16 \times 15 \times 14 = 3,360$)

How many ways can first and second place be awarded to 10 people?

$$\frac{10!}{(10-2)!} = \frac{10!}{8!} = \frac{3,628,800}{40,320} = 90$$

(which is just the same as: $10 \times 9 = 90$)

Notation

Instead of writing the whole formula, people use different notations such as these:

$$P(n, r) = {}^n P_r = {}_n P_r = \frac{n!}{(n-r)!}$$

Example: $P(10,2) = 90$

Combinations

There are also two types of combinations (remember the **order** does **not** matter now):

1. **Repetition is Allowed:** such as coins in your pocket (5,5,5,10,10)
2. **No Repetition:** such as lottery numbers (2,14,15,27,30,33)

1. Combinations with Repetition

Actually, these are the hardest to explain, so I will come back to this later.

2. Combinations without Repetition

This is how lotteries work. The numbers are drawn one at a time, and if you have the lucky numbers (no matter what order) you win!.

The easiest way to explain it is to:

- assume that the order does matter (ie permutations),
- then alter it so the order does **not** matter.

Going back to our pool ball example, let us say that you just want to know which 3 pool balls were chosen, not the order.

We already know that 3 out of 16 gave us: $16 \times 15 \times 14 = 16! / 13! = 3,360$ permutations.

But many of those will be the same to us now, because we don't care what order!

For example, let us say balls 1, 2 and 3 were chosen. These are the possibilities:

Order does matter	Order doesn't matter
1 2 3	
1 3 2	
2 1 3	
2 3 1	1 2 3
3 1 2	
3 2 1	

So, the permutations will have 6 times as many possibilities.

In fact there is an easy way to work out how many ways "1 2 3" could be placed in order, and we have already talked about it. The answer is:

$$3! = 3 \times 2 \times 1 = 6$$

So, all we need to do is adjust our permutations formula to **reduce it** by how many ways the objects could be in order (because we aren't interested in the order any more):

$$\frac{n!}{(n-r)!} \times \frac{1}{r!} = \frac{n!}{r!(n-r)!}$$

That formula is so important it is often just written in big parentheses like this:

$$\frac{n!}{r!(n-r)!} = \binom{n}{r}$$

where n is the number of things to choose from, and you choose r of them
(No repetition, order doesn't matter)

It is often called "n choose r" (such as "16 choose 3")

And is also known as the "Binomial Coefficient"

Notation

As well as the "big parentheses", people also use these notations:

$$C(n, r) = {}^nC_r = {}_nC_r = \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

Example

So, our pool ball example (now without order) is:

$$\frac{16!}{3!(16-3)!} = \frac{16!}{3! \times 13!} = \frac{20,922,789,888,000}{6 \times 6,227,020,800} = 560$$

Or you could do it this way:

$$\frac{16 \times 15 \times 14}{3 \times 2 \times 1} = \frac{3360}{6} = 560$$

So remember, do the permutation, then reduce by a further "r!"

It is interesting to also note how this formula is nice and **symmetrical**:

$$\frac{n!}{r!(n-r)!} = \binom{n}{r} = \binom{n}{n-r}$$

In other words choosing 3 balls out of 16, or choosing 13 balls out of 16 have the same number of combinations.

$$\frac{16!}{3!(16-3)!} = \frac{16!}{13!(16-13)!} = \frac{16!}{3! \times 13!} = 560$$

Pascal's Triangle

You can also use [Pascal's Triangle](#) to find the values. Go down to row "n" (the top row is 0), and then along "r" places and the value there is your answer. Here is an extract showing row 16:

			1	14	91	364	...
		1	15	105	455	1365	...
	1	16	120	560	1820	4368	...

1. Combinations with Repetition



Let us say there are five flavors of icecream: **banana, chocolate, lemon, strawberry and vanilla**. You can have three scoops. How many variations will there be?

Let's use letters for the flavors: {b, c, l, s, v}. Example selections would be

- {c, c, c} (3 scoops of chocolate)
- {b, l, v} (one each of banana, lemon and vanilla)
- {b, v, v} (one of banana, two of vanilla)

(And just to be clear: There are **n=5** things to choose from, and you choose **r=3** of them. Order does not matter, and you **can** repeat!)

Now, I can't describe directly to you how to calculate this, but I can show you a **special technique** that lets you work it out.



Think about the ice cream being in boxes, you could say "move past the first box, then take 3 scoops, then move along 3 more boxes to the end" and you will have 3 scoops of chocolate!

So, it is like you are ordering a robot to get your ice cream, but it doesn't change anything, you still get what you want.

Now you could write this down as $\rightarrow \circ \circ \circ \rightarrow \rightarrow \rightarrow$ (arrow means **move**, circle means **scoop**).

In fact the three examples above would be written like this:

{c, c, c} (3 scoops of chocolate): $\rightarrow \circ \circ \circ \rightarrow \rightarrow \rightarrow$
 {b, l, v} (one each of banana, lemon and vanilla): $\circ \rightarrow \rightarrow \circ \rightarrow \rightarrow \circ$
 {b, v, v} (one of banana, two of vanilla): $\circ \rightarrow \rightarrow \rightarrow \rightarrow \circ \circ$

OK, so **instead of worrying about different flavors**, we have a *simpler* problem to solve: "in how many different ways can you arrange arrows and circles"?

Notice that there are always 3 circles (3 scoops of ice cream) and 4 arrows (you need to move 4 times to go from the 1st to 5th container).

So (being general here) there are $r + (n-1)$ positions, and we want to choose r of them to have circles.

This is like saying "we have $r + (n-1)$ pool balls and want to choose r of them". In other words it is now like the pool balls problem, but with slightly changed numbers. And you would write it like this:

$$\binom{n+r-1}{r} = \frac{(n+r-1)!}{r!(n-1)!}$$

where n is the number of things to choose from, and you choose r of them
(Repetition allowed, order doesn't matter)

Interestingly, we could have looked at the arrows instead of the circles, and we would have then been saying "we have $r + (n-1)$ positions and want to choose $(n-1)$ of them to have arrows", and the answer would be the same ...

$$\binom{n+r-1}{r} = \binom{n+r-1}{n-1} = \frac{(n+r-1)!}{r!(n-1)!}$$

So, what about our example, what is the answer?

$$\frac{(5+3-1)!}{3!(5-1)!} = \frac{7!}{3! \times 4!} = \frac{5040}{6 \times 24} = 35$$